

## WY Implants and Surgical Arts

307-362-4867

Dr. Shaziya Haque, DDS MS

### PATIENT INSTRUCTIONS FOLLOWING IMPLANT PLACEMENT

- **Bleeding:** Small amounts of blood in the saliva can make your saliva appear quite red. This is normal and may be noticed the rest of the day after the procedure.
- **Smoking:** Smoking should be stopped following surgery. Healing and success of the implant will be substantially reduced by the cigarette smoke chemicals in your body.
- **Pain:** Some discomfort is normal after surgery. To minimize pain, take 600 mg of Ibuprofen (three pills) or similar non-aspirin pain reliever every 4-6 hours until bedtime to maintain comfort. If pain medication is prescribed, take it as you need it. Do not exceed the dose on the label. Taking it with food or milk will help reduce upset stomach. Avoid driving or operating heavy machinery when taking pain prescriptions. Do not drink alcohol while taking prescription pain medications.
- **Nausea:** This is most often caused by taking pain medication on an empty stomach. Reduce nausea by preceding each pill with soft food, and a large glass of water.
- **Swelling:** Applying an ice bag to the face over the operated area will minimize swelling. Apply for 15 minutes, then remove for 15 minutes. Continue this for the first day. DO NOT APPLY COLD AFTER 48 HOURS!
- **Numbness:** The local anesthetic will cause you to be numb for several hours after you leave the office. Be very careful not to bite, chew, pinch, or scratch the numb area. Sometimes the implant procedure causes residual numbness or tingling for six weeks or longer.
- **Brushing:** Do not brush your teeth for the first 8 hours after surgery. After this you may brush your teeth gently, but avoid the area of surgery for 2 weeks.
- **Rinsing:** Avoid all rinsing or swishing for 24 hours after your procedure. After 24 hours you may begin gently rinsing with a salt water solution (1/2 teaspoon salt + 1/2 teaspoon baking soda + 8 oz. of water).
- **Diet:** Eat soft foods for the first two days. Maintain a good, balanced diet. Return to normal regular meals as soon as you are able after the first two days. Drink plenty of water. Avoid alcohol for 48 hours.
- **Activity:** After leaving the office, rest and avoid strenuous activities for the remainder of the day. Keeping blood pressure lower will reduce bleeding and aid healing.
- **Antibiotics:** If you were given an antibiotic prescription, take all of them as directed until they are gone. Women, some antibiotics can reduce the effectiveness of birth control pills. Use alternate birth control methods for two months.

- **Sinus:** If your sinus was involved in the procedure you should avoid blowing your nose or playing a wind musical instrument for one week. Use of decongestant medications might be recommended.
- **Removable appliances, dentures:** your dentist will give you specific instructions about your prosthesis. To avoid putting any pressure on the new implants before they have healed, your denture might need to be adjusted or significantly modified. In certain cases, you will need to go without your dentures for a period (days or weeks) after the implants are placed. Sometimes a temporary removable appliance is made for cosmetic purposes, until a new non-removable one can be made.
- **Return** to your dentist office for removal of sutures or follow up checks as directed.

**PLEASE CALL YOUR DENTIST IF YOU HAVE:**

- Uncontrollable pain
- Excessive or severe bleeding
- Marked fever
- Excessive warm swelling occurring a few days after the procedure
- Reactions to medications, especially rash, itching, or breathing problems

**Following these instructions very closely will greatly help your comfort and promote uneventful healing of the area. If any of the instructions are not followed, you might have significantly more discomfort, and the success of the procedure may be affected.**

**Please call us with any questions or concerns.**

**Rock Springs Periodontics: 307-362-4867**

**Dr. Shaziya Haque's Cell: 443-802-5201**